

# Tri-Solfen® Technical Notes



## Tri-Solfen Application Procedures for Cattle

As with many animal health products, ensuring that Tri-Solfen is applied correctly is an important part of the usage process. To achieve the best results, follow these instructions for each of the registered procedures. Registration details are available on the Tri-Solfen packaging label.

Tri-Solfen is a viscous fluid spray, designed for topical use on wounds. Apply Tri-Solfen spray to the wound immediately after the procedure is completed using the Tri-Solfen applicator. It is important that Tri-Solfen is applied evenly as per the instructions below, particularly the wound edges. Do not spray to run off.


## Castration

### DOSE RATES


For castration, each dose of Tri-Solfen is applied in 3 parts. One third inside the vaginal tunic of each testicle and the final third to the edges of the scrotal skin wounds.

Liveweight/Bodyweight	Inside Vaginal Tunic Step 2	External Incisions Step 4	Total Dose
Calves 30kg-100kg	2 x 2ml	1 x 2ml	6ml
Calves over 100kg	2 x 3ml	1 x 3ml	9ml


### APPLICATION




1  
Cut through skin of the scrotum with a scalpel as per normal castration procedure.



2  
Insert the nozzle of the applicator as far into the vaginal tunic as possible and eject 1/3 of the total dose.



3  
Remove testis as per usual procedure. Repeat steps 1–3 on the other side of the scrotum to remove the second testis.



4  
Apply the final third of the dose to the edges of the scrotal incision.

# Tri-Solfen® Technical Notes



## Dehorning and Disbudding

### DOSE RATES

For dehorning and disbudding the dose is applied as 2 x 2ml bands per horn which cross over the wound immediately after each horn is removed.

Liveweight/Bodyweight	Total number of 2ml sprays per animal	Dose per horn	Total Dose
Scoop or knife Dehorning	4	4ml	8ml
Hot Iron disbudding (Calves less than 3 months old)	4	4ml	8ml

### APPLICATION

**Step 1:** Perform disbudding or dehorning for each horn according to accepted procedures.

**Step 2:** Immediately after disbudding or dehorning the first horn, apply Tri-Solfen with the recommended applicator to completely cover the wound including the surrounding burn wound (if using cautery) or the surrounding cut skin edge (if using excision).



### DO NOT SPRAY TO RUN OFF. ENSURE THAT THE PRODUCT FROTHS ON APPLICATION

1. Always use the specially designed Tri-Solfen Applicator for all applications of Tri-Solfen.



CASTRATION NOZZLE

2. When applying Tri-Solfen topically to external wounds always ensure that the applicator is squeezed hard enough to cause the product to froth and cause foaming. This will ensure that the product sticks to the wound and that run off from the application site is minimised.



CORRECT APPLICATION